



References

- Bannerjee S. et al. (2005). Placental expression of interferon-gamma and its receptor IFN-gamma R2 fail to switch from early hypoxic to late normotensive development in preeclampsia. *J. Clin Endocrinol Metab*, *90*(2), 944-952.
- Berczi, I. & Szentvanyi, A. (2003). The Acute Phase Response. In Berczi I and Szentivanyi A, (Eds.), *NeuroImmune Biology: The Immune-Neuroendocrine Circuitry: History and Progress*. (p. 476). New York, NY: Elsevier.
- Dahlgren, J. (2001). Prenatal cytokine exposure results in obesity and gender-specific programming. *Am J Endocrinol Metab*. *281*, E326-334.
- Dodic, M. et al. (2002). Programmed hypertension: kidney, brain or both? *Trends Endocrinol Metab*, *13*(9), 403-408.
- Gabry, K.E. et al. (2003). The Hypothalamic-Pituitary-Adrenal (HPA) Axis: A Major Mediator of the Adaptive Responses to Stress. In Berczi I and Szentivanyi A, (Eds.), *NeuroImmune Biology: The Immune-Neuroendocrine Circuitry: History and Progress*. (pp. 379-414). New York, NY: Elsevier.
- Matthews, S.G. (2002). Early Programming of the hypothalamo-pituitary-adrenal axis. *Trends Endocrinol Metab*, *13*(9), 373-380.

*rats

