



References

- Barker, D.J. (2002). Fetal programming of coronary heart disease. *Trends Endocrinol Metab.* 13(9), 364-368.
- Bessinger, R.C. & McMurray, R.G. (2003). Substrate Utilization and Hormonal Responses to Exercise in Pregnancy. *Clin Obstet Gynecol*, 46(2), 467-478.
- Clapp, J.F. (2004). Maternal insulin-like growth factor-I levels reflect placental mass and neonatal fat mass. *AJOG*, 190(3), 730-6.
- Davies, M.J. & Norman, R.J. (2002). Programming and reproductive functioning. *Trends Endocrinol Metab.* 13(9), 386-392.
- Dodic, M. et al. (2002). Programmed hypertension: kidney, brain or both? *Trends Endocrinol Metab.* 13(9), 403-408.
- Fant, M.E. (2001). Insulin and insulin-like growth factors in Human Development: Implications for the Perinatal Period. *Sem Perinat*, 25(6), 426-435.
- Hesselmar, B. et al. (2002). Born small for gestational age: relation to future allergy and asthma. *Acta Paediatr*, 91, 992-994.
- Holt, R.I. (2002). Fetal Programming of the growth hormone – insulin-like growth factor axis. *Trends Endo Metab*, 13(9), 392-397.
- Kenneway, D.J. (2002). Programming of fetal suprachiasmatic nucleus and subsequent adult rhythmicity. *Trends Endo Metab*, 13(9), 398-402.
- Malek-Khosravi, S. et al. (2004). Insulin changes in preeclamptic women during pregnancy. *Ann Saudi Med*, 24(6), 434-436.
- Ozanne, S.E. & Hales, C.N. (2002). Early Programming of Glucose-Insulin Metabolism. *Trends Endo Metab*, 13(9), 368-373.
- Popovici, R.M. et al. (2001). Hypoxia regulates insulin-like growth factor-binding protein in human fetal hepatocytes in primary culture: Suggestive Mechanism for *in utero* fetal growth restriction caused by uteroplacental insufficiency. *J. Clin Endocrinol Metab*, 80(6), 2653-9.
- Weinstock, M. (2001). Alterations induced by gestational stress in brain morphology and behavior of the offspring. *Progress in Neurobiology*; 65, 427-451.

*rats

